

# PLUMBAGINÆ EUROPÆÆ.

Auctore VICTORE de JANKA.

- I. Scapi simplices monocephali, sub capitulo vagina subulata reversa involuti; spiculæ 1-bracteatae, styli triente inferiore barbati vel plumosi (usque ad basin 5-partiti):

## ARMERIA.

1. «*Macrocentron*: calyx pedicello oblique insertus, infra insertionem productus in calcar attenuato-subulatum hirsutum pedicello contiguum et tubi longitudinem dimidiam saltem æquans, latere interiori pedicelli contactu foveola anguste linearis longa excavatus» (Boissier in DC. Prodr. XII p. 674). 2  
 «*Plagiobasis*: calycis pedicello oblique inserti basis plus minus oblique truncata, latere interiori pedicelli contactu foveola brevi oblonga, ovata vel rotunda excavata» (Boiss l. c. p. 677). 13
2. Phylla (involucri) externa interiora æquantia acuminato-cuspidata vel aristata 3  
 Phylla vel sensim aucta vel exteriora dimidio breviora, pleraque obtusissima 5
3. Folia angusta, linearis-lanceolata ad summum, costæ tubi calycini omnes 10 pilosæ; species basi suffrutescentes 4  
 Folia latiora; costæ calycinae 5 primariae solum pilosæ; haud suffrutescens: *Armeria gaditana* Boiss.
4. Caules dense foliati, parte nuda vix exserti; limbus calycinus vix distincte lobatus erosus, abrupte mucronulatus; tota planta velutino-tomentosa: *A. cinerea* Boiss. et Welw.  
 Caules nudi folia superantes; limbi calycini lobi 3-angulares mucronulati; planta glabra: *A. Welwitschii* Boiss.
5. Limbi calycini lobi obsoleti vel indistincti, mucronati vel mutici 6  
 Calycis lobi distincti. 7.
6. Folia utplurimum elongata anguste linearia vel setacea (6—10 poll. longa, lineam lata) 1-nervia; bracteæ interflorales obsoletæ:  
*A. macrophylla* Boiss. et Reut.

- Folia abbreviata oblongo-lanceolata (2—4 poll. longa,  
2—3 lineas lata) plurinervia; bracteæ interflorales calyces inte-  
riores æquantes --- A. *velutina* Welw.
7. Limbus calycinus tubo æquilogus vel longior --- 8  
Limbus calycinus tubi dimidiata vix superans --- 12
8. Folia scapo semper angustiora plus minusve arcuata vel flexuoso-  
curvula; species suffrutescentes --- 9  
Folia latitudine scapi vel latiora (3-nervia) --- 10
9. Folia hirtula linearis-setacea canaliculato 3-queatra elongata; scapi  
elati; involucrum coriaceum velutinum --- A. *pinifolia* R. et Sch.  
Folia glabra subcompressa abbreviata flexuoso-curvula;  
scapi humiles semper arcuato-adscendentibus; involucrum omnino  
scariosum --- A. *arcuata* Welw.
10. Bracteæ interflorales deficientes; involuci phylla pallida anguste  
marginata [aristæ lobis calycinis breviores; folia (2-formia:  
primaria latiora) flaccida] --- A. *Boissieriana* Coss.  
Bracteæ interflorales ± evolutæ; folia haud flaccida --- 11
11. Bracteolæ interflorales minutæ pedicellos fructiferos vix  
superantes; lobi calycini validiuscule (breviter) mucronati (folia  
rigida) --- A. *baetica* Boiss.  
Bracteolæ interflorales totum calyceum subæquantes; lobi  
calycini tenuiter longe aristati --- A. *rumelica* Boiss.
12. Bracteolæ interflorales fere omnino deficientes; scapi graciles  
. A. *pungens* R. et Sch.  
Bracteolæ interflorales minimæ, tamen fere pedicellorum  
fructiferorum longitudine; scapi fistulosi: --- A. *fasciculata* Willd.
13. Folia subjuniperino-triquetra columnari-imbricata; scapi e  
cæspite vix exserti; capitula subsessilia («limbus calycinus e  
sinubus aristatus») --- A. *cæspitosa* Boiss.  
Folia elongata haud aceroso-triquetra; scapi elatiore --- 14
14. Tubi calycini costæ 5 principales solum pilosæ, reliquæ inter-  
vallaque glaberrime; folia conformia lata, lanceolata; (calycis  
limbus subtruncatus, lobi latissimi brevissimi, aristæ brevissi-  
mæ mucroniformes, crassiusculæ, incurvæ) --- A. *santa* Janka.\*

\* E rupibus maritimis pr. coenobium Lavra ad ped. m. Athos. — Specimina a  
celeberr. Boissier in Fl. or. IV. pag. 873 infauste ad var. brachiphyllam Armeriae majel-  
lensis Ten. — A Orphanidis Boiss. relata. — Planta athoa foliis semper uniformibus atque  
præcipue costis calycinis 5 solum pilosis et aristis brevissimis introrsum curvis distinc-  
tissima! Quoad calycem ad costas primarias tantum pilosum cum A. cariensi Boiss.  
convenit, quæ vero ob cicatricem linearē ad aliam sectionem pertinet. — Habitu nova  
species plane A. Gussonei similis.

Costæ calycinæ omnes 10 pilosæ	15
15. Involueri phylla 2—3 infima cæteris longiora totumque florū capitulum in alabastro juvenili longiuscule superantia	16
Involueri phylla exteriora capitulum haud, vel rarissime breviter excedentia	17
16. Folia linearis-lanceolata	<i>A. plantaginea</i> Willd.
Folia linearia	<i>A. elongata</i> Willd.
17. Costæ calycinæ secundariæ indistincte, saltem parcius vel obsolete pilosæ (flores albi).	18
Costæ omnes 10 æqualiter pilosæ	19
18. Folia conformia (latiuscule linearis-lanceolata); phylla involucrantia fere ex toto herbacea	<i>A. pubinervis</i> Boiss.
Folia difformia [exteriora (primaria) lanceolata vel lanceolato-linearia, — interiora linearia v. linearis-setacea]; phylla involucralia bracteæque niveo-splendentes	<i>A. argyrocephala</i> Wallr.
	( <i>A. undulata</i> Boiss.)
19. Phylla involucralia exteriora longe attenuato-acuminata, cuspido-aristata vel pungenti-mucronata	20
Phylla involucralia exteriora acutiuscula brevius cuspidata vel mutica	22
20. Lobi calycini obsolete (ergo limbus calycinus subtruncatus); folia glaberrima	<i>A. latifolia</i> Willd.
Lobi calycini elongati; folia ± pilosa vel ciliata (flores albi)	21
21. Phylla involucralia subpungenti-mucronata; folia 2-formia (vetusta vel primaria novellis latiora)	<i>A. villosa</i> Gir.
Phylla involucralia molliter i. e. inermi-aristata, folia conformia	<i>A. longearistata</i> Boiss
22. Longearistatae: aristæ lobos calycinis æquantes vel superantes	23
Breviaristatae: aristæ lobis calycinis breviores	32
23. Folia brevia latave latitudine 3-plo circiter longiora	<i>A. Morisii</i> Boiss.
Folia elongata angustiora	24
24. Folia omnia conformia	25
Folia primaria novellis latiora	29
25. Folia lineaearia vel anguste linearis-lanceolata	26
Folia latiora	28
26. Involueri phylla subæquilonga pallida; folia acuta vel mucronata; flores albi	27
Involueri phylla («exteriora dimidio breviora») bracteæque rubellæ vel purpurascentes; folia obtusata; flores purpurascentes	<i>A. alpina</i> Willd.

27. Bracteolæ interflorales valde elongatae phylla involucralia longe superantes; folia recta: --- A. *leucocephala* Koch.  
Bracteolæ interflorales haud ita elongatae; folia curvata  
A. *ruscinonensis* Gir.

28. Limbi calycini tubo æquilongi truncati lobi brevissime 3-angulares (flores albi) --- A. *allioides* Boiss.  
--- A. *bupleuroides* Godr. et Gren.  
Limbi tubo longioris lobi ovati --- A. *macropoda* Boiss.

29. Folia obtusa («phylla involucralia fere æquilonga»); limbi calycini nervi versus medium loborum evanidi --- A. *majellensis* Boiss.  
Folia acuta --- 30

30. Phylla involucralia interna late membranacea externis duplo longiora; limbus calycinus tubum æquans --- 31  
Phylla involucralia sensim aucta anguste membranacea;  
limbus calycinus tubo brevior (folia 3-nervia) --- A. *canescens* Boiss.\*

31. Folia 1-nervia --- A. *nebrodensis* Guss.  
Folia 3-nervia --- A. *Orphanidis* Boiss.

32. Folia lata, oblongo-lanceolata vel lanceolato-linearia --- 33  
Folia anguste linearia --- 34

33. Phylla involucralia scariosa; folia 5—7-nervia --- A. *Gussonei* Boiss.  
Phylla involucri coriacea; folia 3-nervia --- A. *cantabrica* Boiss.

34. Folia difformia --- 35  
Folia conformia --- 39

35. Phylla involucralia omnia æquilonga vel subæquilonga --- 36  
Phylla involucri sensim aucta --- 37

36. Phylla involucralia dorso herbacea pleraque acutissima; lobi calycini ob limbum subtruncatum obsoleti abrupte aristulati  
--- A. *denticulata* Bert.  
Phylla involucralia scarioso-membranacea, anteriora obtusissima; lobi calycini distincte ovati --- A. *juncea* Gir.

37. Phylla involucri omnia dorso excurrente + longe mucronata;  
limbus calycis tubum æquans --- A. *Durieui* Boiss.  
Phylla interiora obtusa vel retusa; limbus calycinus tubo brevior --- 38

\* Species etiam in Italia indigena: Armeria floribus roseis a me in provinc. Lucania Ital. inferioris, in Apennini cacuminibus lapidosis (inter Laviano et Muro d. 22. Jun. 1874 m. Eremito) lecta certe ad A. canescentem Boiss. pertinet. — Corolla quidem «constanter alba» (cfr. Boiss. in DC. Prodr. XII. p. 686) debet esse, tamen roseas video in ipsis speciminiibus dalmaticis Pichlerianis 1870 lectis. — Eandem exakte plantam e Calabria rite ad speciem dalmaticam retulerunt jam a 1877 Huter, Porta et Rigo sec. schedulam collectorum ex itinere italico III n. 305 b «e Calabriae oriental. distr. Castrovillari pasc. silvat. 18. Jun. mte Cotorrocole».

Scapi vel caules ramosi; spiculae 3-bracteatae; styli glabri  
vel papilloso-asperi

II

- II. *Caulis elatus foliis plerisque ample auriculato-amplexicaulibus  
alte vestitus; flores ad ramorum apicem subcapitato-spicati; co-  
rollæ hypocraterimorphæ tubus calycem longe superans; stylus  
simplex, dimidio inferiore papilloso-hirtulus **PLUMBAGO europaea L.***

Scapus, vel caulis solum squamatus, rarius basin versus foliatus vel altius, sed tunc folia semper basi attenuata, haud ita auriculata; *stylus ad medium vel profundius divisus* — —

三

- III.** Folia brevia angustave acerosa, plerumque juniperina subulato-triquetra, rarius planiuscula; (suffruticulus cæspitosus erinaceus: scapi vix e cæspite exserti apice spicas brevissimas usque 8-spiculatas gerentes); styli basi inter se coalitæ lèves, *stigmata subcapitata*; bracteæ indivisæ — **ACANTHOLIMON Echinus** Boiss.

Folia nunquam acerosa, latiora; scapi vel caules elatiores IV

IV

- IV.** Ob bracteam intermedium minutam linearis-triquetram infima fere omnino occultam solum duæ exteriore amplæ spectabiles; *styli ad medium usque connati* glabri, stigmatibus filiforme-cylindricis (corolla usque ad faucem gamopetala, caulis alte, rami ramulique crebre foliati) — **LIMONIASTRUM monopetalum L.**

Bracteæ tres sursum majores distinctissimæ ; scapi aphylli vel caules ima basi confertius foliati ; styli omnino vel usque ad imam basin liberi

V

- V. Bracteæ interiores 3-cuspidatae, vel præter cuspides 1—2 terminales lateraliter angulato-auriculatae; *styli tota longitudine papilloso-asperi*; *stigmata capitata*:

## GONIOLIMON.

1. Humile, vix 2-pollicare, foliorum lamina 3—4 lin. longa, panicula pollicaris, spiculæ paucæ fasciculatae; bracteæ excurvæ; calycis tubus obsolete glandulosus ... *Goniolimon Sartorii Boiss.*  
Majora ... 2

2. Spiculæ (2—3) fasciculatae; tubus calycinus glaber demum dimidio superiore e bracteis exsertus ... 3  
Spiculæ spicatae; tubus calycinus ad costas + villosulus bracteis altius involutus ... 4

3. Spicularum fasciculi confertiusculi; corollæ limbo calycino concolores albæ! ... *Goniolimon collinum Boiss.*  
(*G. dalmaticum* Boiss. Fl. orient. — minime Rehb. fil.)  
Spicularum fasciculi inter se remoti; corollæ limbo calycino discolores persicinæ! ... *G. Besserianum Nyman.*  
(*Statice Besseriana* R. et Sch.)

4. Spiculæ linearis-picatae, rami apice scorpioidei flexiles ... 5  
Spicæ abbreviatæ ovales; rami effusi divaricato-patuli rigidissimi bracteisque horridi ... *G. dalmaticum Rehb. fil.*  
(*Statice dalmatica* Presl.)

5. Bracteæ intimæ cuspides 2 tantum evolutæ, tertia (lateralis) ad dentem 3-angulum nembranaceo-marginatum reducta; sinus inter cuspides membranaceæ: ... *G. Beckerianum Janka.*  
(*Statice incana* Becker pl. exs. e Sarepta a l Wolgam inferiorem — non L.)  
Bractea intima semper 3-cuspidata; sinus inter cuspides herbaceaæ ... *G. tataricum Boiss.*  
(*Statice tatarica* L.)

Bracteæ indivisæ vel rarius intima ad summum repando-emarginata: *styli* glabri; *stigmata* filiformi-cylindracea:

## STATICE :

Inflorescentiae rami haud ita dense squamati; folia basilaria latiora, haud linearia plerumque persistentia	5
4. Flores in ramorum apicibus densissime corymboso-conferti paniculam myriantham formantes; squamulae tenuissime elongate aristato-caudatae, quasi pilo terminatae	<i>St. ferulacea L.</i>
Flores versus ramorum extremitates laxe racemoso-spicati axi communi ipso apice sterili superati; squamulae breviter aristatae	<i>St. diffusa Pourr.</i>
5. Florum arce imbricatorum glomeruli vel capitula interrupte spicati	<i>St. suffruticosa L.</i>
Inflorescentia haud racemoso-spicata	6
6. Petala usque ad limbum in tubum elongatum calycem manifeste excedentem laxe connata. — Herbae fragillimae, ramis inferioribus sterilibus numerosis, corollis pulchre roseis post anthesin non corculatis, calycis cylindraceo-tubulosi vel tubuloso-infundibuliformis limbo omnino exerto bracteam interiore duplo vel fere duplo superante	7
Petala ima basi solum connata, ceterum libera, calyx bracteam intimam minus excedens	8
7. Paniculae amplae ramosissimae ramuli graciles scorpioidei; spiculae graciles, angustae florae numerose spicatae; tubus calycinus glaber	<i>St. cæsia Girard.</i>
Paniculae subpyramidatae rami validiusculi erecti; spiculae robustae plerumque 2-florae in ramulis paucæ; tubus calycinus puberulus	<i>St. insignis Coss</i>
8. Limbi calycini sinus denticuliferi; limbus ideo quasi inæqualiter 10-lobus	9
Dentes inter lobos calycinos nulli; fere omnes palminerviae	14
9. Plantahumilis, gracilis, folia pollicie breviora vel parum longiora, lanceolata, acuta spinula tenui 2—3 lineas longa terminata margine crispatula, scapi squamæ atque ramorum ramulorumque spinuloso-mucronatae	<i>St. spinulosa Janka</i>
(St. spec. nova e ditione Flumineusi herb. Sadler n. 10804 mus. nat. hungar.)	
Elatiores vel robustiores, haud ita spinulosæ, foliis penninerviis	10
10. Lobi calycini acuti vel acutiusculi	11
Calycis lobi rotundati obtusissimi	13
11. Scapus angulatus; spiculae valde inter se dissitae (bracteæ inferiores fere æquilongæ, suprema 2-plo longior; calyx undique pilosus)	<i>St. bahusiensis Fries.</i>
Scapus teres; spiculae approximatæ vel confertæ; calyx ad costas 2 pilosus	12

12. Spiculæ in panicula densa stricta subcorymbosæ ; bracteæ 2 inferiores subæquilongæ, suprema 2-plo longior ... *St. Limonium L.*  
 Paniculæ laxæ haud corymbosæ rami patuli recurvi; bractea infima secunda fere 2-plo, suprema 3-plo brevior ... *St. serotina Rchb.*

13. Tomentella ; bractea infima ovato-3-angularis cuspidata dorso herbacea ... *St. tomentella Boiss.*  
 Glabra; bractea infima ovato-rotunda fere omnino membranacea ... *St. Gmelini Willd.*

14. Rami steriles nulli vel pauci, simplices vel parum ramosi, semper arrecti vel stricte ramosi ... *St. Gmelini Willd.* 15  
 Rami steriles numerosi, articulato - multifidi, utplurimum flexuoso-divaricato-ramosissimi ... *St. Gmelini Willd.* 42

15. Spiculæ (saltet pleræque) subinordinate corymbuloso-fasciellatæ divergentes vel divaricatae ... *St. corymbulosa Coss.* 16  
 Spiculæ regulariter (pectinatum) parallele ordinatae ... *St. Sibthorpiana Guss.* 18

16. Limbus calycinus tubo subæquilongus vel longior ... *St. corymbulosa Coss.* 17  
 Limbus calycinus tubo multo brevior ; (folia ampla, bractea superior infima 3-plo longior ; spiculæ 2—5-floræ) *St. ovalifolia Girard.*

17. Spiculæ 1-floræ ; bractea superior infima 4-plo longior ; lobi calycini nervis validis percorsi ; tubus ad costas longiuscule patuleve hirsutus (folia parva semper mucronata, obovato-spathulata) ... *St. Sibthorpiana Guss.*  
 Spiculæ 2—3-floræ ; bractea superior infima 6-plo longior ; limbi calycini nervi longe ante apicem evanidi ; tubus ad costas pilosus (folia ampla) ... *St. ovalifolia Girard.*

18. Spiculæ plerumque approximatissimæ eximie regularissime distichæ imbricatae (rarissime : in sola St. occidentali laxis dispositæ, sed tunc paniculæ rami strictissimi !) ... *St. Dufourei Girard.* 19  
 Spiculæ haud talimodo distichæ nec ita compactæ ... *St. Girardiana Guss.* 24

19. Spiculæ approximatissimæ spicas crassas compactas abbreviatas formantes ; bractæ obtusissimæ ; spiculæ eximie regularissime distichæ imbricatae ... *St. Girardiana Guss.* 20  
 Spicæ + elongatae lineares vel sublineares minus compactæ ; bractæ + acutæ ... *St. Girardiana Guss.* 21

20. Tota planta (præter folia) pilis brevibus subfasciculatis albis tomentella (rami inferiores steriles) ; spicæ ovato-oblongæ ; bractea superior cuneato-obovata subretusa ; paniculæ rami stricti ... *St. Dufourei Girard.*  
 Glabra ; rami (steriles nulli) patuli ; bractea superior sub apice obsolete mucronulata (folia parva ovata acuta longiuscule petiolata) ... *St. Girardiana Guss.*

21. Rami strictissimi (spicae lineares) ... *St. Girardiana Guss.* 22

Rami patuli vel excurvi recurvique	23
22. Spicæ crassæ robustæ; bracteæ albo-marginatæ; spiculæ eximie distiche dense imbricatae; antheræ sublineares (folia obovato-spathulata obtusissima mutica vel apice brevissime mucronata)	
St. Dodartii Girard.	
Spicæ tenues graciles; bracteæ (superior exteriore i. e infima 2-plo longior) nitide rubello-marginatae; spiculæ minus regulariter laxiusque imbricatae; antheræ ovatae (folia lanceolato-spathulata acutiuscula longiuscule setaceo-apiculata)	
St. occidentalis Lloyd.	
23. Bractea infima acutiuscula; paniculæ rami patuli	
St. lychnidifolia Girard.	
Bracteæ acutæ (brunneæ, rufesceni-marginatae); spicæ strictiusculæ	
St. densiflora Guss. (St. oxylepis Boiss.)	
24. Lobi calycini nervis validis rufescentibus limbum cæterum + perdurantibus atque indurantibus percursi	25
Lobi calycini nervi haud indurantes	27
25. Calycis tubus spinis 5 apice hamatis (i. e. nervis cætero limbo cito emarcido persistentibus valde induratis) patulis terminatus (spiculæ graciles longiuscule cylindraceæ, curvulae, inter se valde dissitæ; herba tenella annua)	
St. echooides L.	
Limbus calycinus minus fugax, nervique calycini minus rigidi nec ita manifeste hamati; spiculæ breviores; herbæ perennes	26
26. Limbi calycini nervi usque ad apicem loborum producti, apice uncinato-inflexi; bractea infima suprema 3-plo brevior:	
St. Companyonis Gren.	
Limbæ calycini nervi paulo ante apicem loborum evanidi apice recti — haud uncinatuli; bractea infima suprema 5-plo brevior	
St. duriuscula Girard.	
27. Bracteæ superiores acutæ vel mucronulatae (folia margine revoluta)	28
Bracteæ superiores obtusissimæ	30
28. Paniculæ rami gracillimi patenti-recurvi; folia majuscula obovato-spathulata	
St. psiloclada Boiss. (St. panormitana Todaro.)	
Paniculæ rami strictiusculi firmiores; folia parva	29
29. Bractea suprema infima apice patula 3-plo longior; limbus calycinus tubum æquans, lobis profundis lanceolatis acutis; folia oblongo-spathulata apica retusa vel emarginata	
St. minuta L.	
Bractea suprema infima adpressa 5-plo longior; limbus	

- calycinus tubo dimidio brevior, lobis oblongis obtusis; folia angustissima, linearis-oblonga, obtusa vel acuta (scapi humillimi, panicula depauperata divaricata!) *St. rupicola Badarro.*
30. Folia amplissima penninervia; herba tota molliter velutina, bracteæ fere omnino albo-hyalinæ; panicula ampla dissitiflora *St. latifolia L.*  
 Folia palminervia parva; indumentum nullum; bracteæ magis herbaceæ *31*
31. Bractea suprema infima 5-plo longior (spiculæ in paniculæ diffusæ ramis ramulisque gracillimis arcuato-excurvis manifeste dissitæ!) *St. salsuginosa Boiss.*  
 Bractea suprema infima 2—3-plo longior *32*
32. Spiculæ valde inter se dissitæ, rami simplices vel subsimplices pauciflori; bractea suprema arcte involuta: spiculæ cylindricæ, graciles *33*  
 Spiculæ (pleræque saltem) ± imbricatim confertæ v. subconfertæ *34*
33. Spiculæ maximæ, fere semipollulares; bracteæ teneræ late membranaceæ (obtusissimæ; folia carnosa cuneato-spathulata in petiolum longe attenuata apice acute emarginata; caudex fruticulosus lignescens) *St. emarginata Willd.*  
 Spiculæ multo minores; bracteæ coriaceæ anguste membranaceæ *St. Sieberi Boiss* *35*
34. Folia plana *35*  
 Folia margine revoluta *40*
35. Bracteæ amplæ glumaceæ valde coriaceæ laxæve, sursum dilatatae spiculas sursum tumidas, inverse conicas vel subelavatas formantes, obscure membranaceo-marginatae; limbus calycinus abbreviatus *36*  
 Bracteæ teneriores haud glumaceæ *37*
36. Folia obovato-spathulata; calycis tubus totus pilosus; spicæ robustæ, crassæ *St. ocytifolia Poir.*  
 Folia oblongo-spathulata; calycis tubus basi tantum pilosus; spicæ graciles; spiculæ (antecedente) dimidio minores:  
*St. corinthiaca B. et H.*
37. Spiculæ brevissimæ; limbus calycinus e bractea parum exsertus tubo brevior *St. delicatula Girard.*  
 Spiculæ longiores; limbus calycinus tubo longior magis manifeste exsertus *38*
38. Bractea infima suprema 2-plo brevior «lanceolata» acuta (v. «acuminata»); limbi calycini lobi lanceolati acutiuseuli; «panicula lata scapo reliquo longior» (bracteæ late membranaceæ)  
*St. globulariæfolia Desf.*

- Bractea infima suprema 3-plo brevior ovalis obtusiuscula  
vel obtusa!; limbi calycini lobi ovato-oblongi otbusi ... ... 39
39. Bracteæ late albo-membranaceo-(scarioso-) marginatæ : margo  
cum parte herbacea haud confluua (spiculæ ob bracteam supre-  
mam valde ventricosam laxam, ovatæ) ... ... ... St. Costæ Willk.
- Bracteæ obscure vel obsoletius rufescenti-membranaceo-  
marginatæ (spiculæ ob bracteam supremam valde involutam  
i. e. calycem valde involventem cylindraceæ; «panicula oblonga  
scapo reliquo brevior confertiflora») ... ... ... St. confusa G. et G.  
(St. Legrandi Timb. et Gaut!)
40. Folia obovato-cuneata ... ... ... ... St. albida Guss.
- Folia angustiora ... ... ... ... 41
41. «Bracteæ duæ exteriores obtusissimæ et fere emarginatæ» (sic  
apud Gussone Fl. siculæ synops. I pag 570; — specimina Parla-  
toreana et a cl. Todaro accepta bracteas exteriores acutas  
habent); folia oblongo-spathulata ... ... ... St. minutiflora Guss.
- Bracteæ acutæ!; folia lineari-oblonga ... St. inarimensis Guss.
42. Bracteæ infimæ omnino hyalino-membranaceæ ... ... ... 43
- Bracteæ infimæ latius herbaceæ ... ... ... 45
43. Bractea inferior subrotunda suprema 4-plo brevior St. dichotoma Cav.
- Bractea inferior angustior superiore 2—3-plo brevior ... 44
44. Bractea inferior obtusa; paniculæ rami divaricati; spiculæ  
breviter conferteque glomerato-spicatae ... ... ... St. caspia Willd.
- Bractea infima acuminata; paniculæ rami (valde elongati)  
erecto-patuli; spiculæ remotaæ laxe longiusque spicatae St. Dubyei G. et G.
45. Glaberrimæ læves; rami steriles stricte ramulosi ... ... 46
- ± Scabridæ, tuberculato-asperæ vel pubescentes ... ... 48
46. Ramulorum articuli nullibi constricti, æqualiter cylindracei ... 47
- Ramuli ad articulationes constricti fere strangulati:  
St. cosyreensis Guss.
47. Paniculæ rami patulo-recurvi; spiculæ cylindraceæ; bractea  
infima acuta ... ... ... ... St. virgata Willd.
- Paniculæ rami stricti; spiculæ ob bracteas supremas  
ventricosas (subinflatæ) ovales (insuper breviores quam in  
St. virgata!); bractea infima obtusa! ... ... ... St. catanensis Tin.  
(Todaro fl. sic, exs. 379!)
48. Scabridæ vel tuberculato-exasperatæ ... ... ... ... 49
- Velutino-puberulæ vel tomentosæ ... ... ... ... 53
49. Ramulorum articuli omnes continue contigui, cylindracei, herbæ  
minus fragiles ... ... ... ... ... ... ... 50
- Articuli ad extremitates repentine constricti i. e. isthmi  
strangulati, stipite brevissimo tenuissimoque quidem sed sat

- manifesto inter se connexi; supremi ovati vel subglobosi;  
 spiculæ curvæ; tubus calycinus ad angulos vel basi  $\pm$  pilosus;  
 herba fragillima ... ... ... ... ... ... *St. articulata Lois.*
50. Calyces bractea obvallati sessiles ... ... ... ... ... ... 51  
 Calyces bractea patula denudati distinctissime pedicellati  
*St. Calcaræ Todaro fl. sic. exs. 276!*  
 (St. dichotoma Guss. Synops.)
51. Spiculæ gracillimæ (rectæ) *St. dubia Andrz.* Todaro fl. sic. exs. 1094!  
 Spiculæ crassiores ... ... ... ... ... ... 52
52. Bractea suprema margine anguste membranacea; spiculae  
 rectæ; tubus calycinus glaber (caules nigrescentes) *St. dictyoclada Boiss.*  
 Bractea suprema usque ad dimidiā submembranacea;  
 tubus calycinus sparse pilosus; spiculæ curvæ ... *St. rorida S. et Sm.*
53. Rami ramulique steriles rectangule infracti flexuoso-divarica-  
 tissimi: bractea suprema infima 4-plo longior; limbus calycinus  
 tubo brevior ... ... ... ... ... ... *St. cancellata Bernh.*  
 Rami nunquam refracti; limbus calycinus tubum æquans 54
54. Laxifloræ; bractea suprema infima 6-plo longior ... ... ... ... ... 55  
 Inflorescentia densa fastigiato-corymbosa; bractæ velu-  
 tinæ, infima albo-marginata suprema 3-plo brevior; folia  
 obcordata ... ... ... ... ... ... *St. pubescens DC.*
55. Bractæ glabræ: infima fere omnino membranacea; folia  
 obcordata; herba breviter furfuraceo-velutina ... ... *St. cumana Ten.*  
 Bractæ velutinæ: infima vix margine membranacea;  
 folia spathulata; herba dense patule velutina ... ... *St. furfuracea Lag.*
-

## BRASSICEÆ EUROPÆÆ.

Auctore VICTORE DE JANKA.

1.	Siliqua valde abbreviata, globosa didyma vel ovoideo-globosa (sepala erecta, petala longe exserte unguiculata) ---	2
	Siliqua elongata ---	6
2.	Herbae annuae, foliis pinnatisectis; filamenta libera ---	3
	Frutices vel suffrutices lignosi rigidi ramosissimi, foliis indi- ysis fere semper integerrimis; filamenta majora per paria connata	4
3.	Siliquæ costis breviter parceve dentato-echinatis, rostro foliaceo subcochleari, pedicello inflexo nutantes (petala violaceo-venosa)	
	CARRICHTERA <i>Vellæ DC.</i>	
	Siliquæ tota superficie longe echinato-hispidissimæ rostro tetragono-subulato, erectæ (petala avenia) <b>SUCCOWIA</b> <i>balearica L.</i>	
4.	Siliculæ setoso-hispidissimæ stylo anguste linguæformi rostratae (racemi abbreviati densiflori; suffrutex elatior inermis; petala fusco-venosa) ---	
	<b>VELLA</b> <i>aspera Pers.</i> ( <i>Boleum asperum Desv.</i> )	
	Siliculæ glabrae rostro late ovato foliaceo ---	5
5.	Fruticulus humilis dense cæspitosus ramiculis superioribus in spi- nas induratas transformatis undique horridus, erinaceus; racemi abbreviati laxe 3—5 flori; folia linear-lanceolata vel linea- ria (petala violaceo-reticulata) ---	
	<i>Vella spinosa Boiss.</i>	
	Suffrutex elatior inermis; flores longe spicato-racemosi; folia obovata vel obovato-lanceolata ---	
	<i>V. Pseudo-Cytisus L.</i>	
6.	Siliqua lanceolato-oblonga, basi attenuata, valvis nervis 5 validis rectis æquidistantibus percursis; <i>semina</i> toto ambitu late membra- naceo-alata; <i>stamina</i> 4 longiora per paria filamentis usque ad apices coalita. Frutex floribus albis, petalis venis fuscis reticulatis	
	<b>EUZOMODENDRON</b> <i>Bourgæanum Coss.</i>	
	Siliqua utrinque æqualiter vel solum apicem versus attenu- ata: <i>semina</i> exalata vel rarius unilateraliter alata; filamenta libera ---	7

## BRASSICA

7.	Folia caulina suprema (indivisa) <i>deorsum in auriculas amplas producta vel profunde cordato-amplexicaulia</i> ...	8
	Folia haud ita profunde auriculata vel cordato-amplexicaulia	14
8.	Pedicelli siliquas superantes (stylus 1''' longus, gracilis, siliquæ torulosæ) ...	<i>Brassica amplexicaulis DC.</i> (sub Sinapide)
	Pedicelli siliquis breviores	9
9.	Flores flavi v. lutei; folia basilaria et caulina inferiora lyratopinnatiflora	10
	Flores purpurascentes, violacei v. albi; folia indivisa.	11
10.	Flores corymbosi i. e. gemmas superantes: ...	<i>Brassica Rapa Koch.</i>
	Flores racemosi i. e. gemmis superati	<i>B. Napus L.</i> ( <i>B. campestris DC.</i> )
11.	Folia omnia obtusa; siliquæ plano-compressæ; semina 2-seriata, rostrum 2''' longum	<i>B. Moricandia Boiss.</i> ( <i>Brassica arvensis L.</i> , <i>Moricandia arvensis DC.</i> )
	Folia superiora acuta; siliquæ tetragono-compressæ vel teretiusculæ; semina 1-seriata, rostrum brevius	12
12.	Pedicelli crassi breves; rostrum breve conicum; petala violaceo-purpurea	13
	Pedicelli graciles 5—6''' longi; rostrum linearis-conicum; petala alba (parva)	<i>B. fætida Bourg.</i> (sub <i>Moricandia</i> )
13.	Siliquæ compresso-tetragonæ venis distincte prominulis; flores magni, semina anguste albo-marginata	<i>B. moricandoides Boiss.</i> ( <i>Moricandia Ramburei Webb.</i> )
	Siliquæ teretiusculæ obsolete venosæ; flores parvi, semina (minora) latius marginata:	<i>B. anticaria Rouy in litt.</i> ( <i>Moricandia baetica B. et K.</i> )
14.	Folia caulina suprema (indivisa) basi dilatata semiamplexicaulia	15
	Folia caulina suprema basi angustata sessilia vel petiolata, nunc caulis omnino aphyllus	23
15.	Folia vel tota herba indumento molli vestita	16
	Glaberrimæ vel sparsissime setulosæ	19
16.	Flores parvi	17
	Flores magni	18
17.	Siliquæ compressæ sub-6-spermæ, sesquipollicares; stylus <i>ovato-lanceolatus</i> siliquæ diametr. longus i. e. $1\frac{1}{2}$ linearis aspermus	<i>B. Botteri Vis.</i>
	Siliquæ 4-gonæ, 12—16-spermæ; stylus conico-subulatus lineam longus	<i>B. mollis Vis.</i>

18. Folia tomentoso-incana ; siliquæ teretes elongatae ( $2\frac{1}{2}$ —3 pollicares) graciles substipitatae ; stylus 3—4 lin. longus, tenuis, subulatus 1—2-spermus (nonnisi abortu 1-spermus ; siliquæ ut in *B. oleracea*) ; corolla alba ... ... ... ... *B. incana* Ten.  
     Folia villosa-canescens ; siliquæ sub-4-gonæ pollicem haud excedentes, non stipitatae ; stylus cylindricus 1''' longus aspermus ; corolla flava ... ... ... ... *B. villosa* Biv.
19. Siliqua abbreviata 1— $1\frac{1}{2}$  pollicaris, *crassissima* (fere Raphani sativi !) ; stylus  $\frac{1}{2}$  vel  $\frac{1}{3}$  siliquæ metiens : ... ... *B. macrocarpa* Guss.  
     Siliqua longior et gracilior, stylus brevior ... ... ... ... 20
20. Inferne suffrutescentes, ramis basi lignescens ... ... ... ... 21  
     Herbaceæ annuae vel biennes ... ... ... ... 22
21. Flores candidi ... ... ... ... *B. nivea* Boiss. et Sprun.  
     Flores flavi (folia numquam lyrata, suprema integerrima) *B. cretica* L.
22. Folia superiora incisa ; siliquæ teretiusculæ ... ... ... ... *B. oleracea* L.  
     Folia superiora integra ; siliquæ tetragonæ ... ... ... ... *B. montana* Pourr.
23. Siliquæ rostrum nunc teres, paulo (saepius) tetragono-compressum, nunc totum (siliquæ tunc gracile linearis valvis multo brevius) vel solummodo apice ancipiti-compressum ... ... ... ... 24  
     Siliquæ rostrum totum plano-compressum, foliaceum, anceps, ensiforme-elongatum siliquæ breviusculæ atque crassioris dimidiadim æquans vel longius ... ... ... ... 67
24. Siliquæ rachi adpresso ... ... ... ... 25  
     Siliquæ rachi haud adpresso in pedicellis patulis vel patentissimis ... ... ... ... 27
25. Stylus vix lineam longus ... ... ... ... 26  
     Stylus 3''' vel ultra longus ; (pedicelli calycem superantes)  
         *B. pubescens* Ardoino  
         (Flore des Alpes maritimes ; — *Sinapis pubescens* L.)
26. Siliquæ angustæ, tenues, cylindraceo-subulatae valde torulosæ ; sepala pedicellis breviora patentissima ; pedicelli crasse cuneati ... ... ... ... *B. adpressa* Boiss.  
         (Sinapis incana L.)  
     Siliquæ latæ, conico-attenuatæ, crassæ ; sepala pedicellis longiora patula ; pedicelli tenues ... ... ... ... *B. nigra* Koch.  
         (Sinapis nigra L.)
27. Inflorescentia foliaceo-bracteata i. e. folia sensim insensibiliter decrescentia per inflorescentiam continua atque pedicellos inferiores saltē stipantia ... ... *B. bracteata* Gren. et Godr. (sub *Diplotaxide*)  
         (B. Pollichii Jess. — *Erucastrum Pollichii* Sch. et Sp.)  
     Folia interfloralia nulla ... ... ... ... 28
28. Pedicelli longissimi, siliquas pluries superantes (folia pectinato-

pinnatipartita, lobis acutis, lateralibus semi-ovatis margine inferiore integerrimis, superiore inciso-dentatis	... ... ...	<i>B. Barrelieri DC.</i>
		(sub <i>Diplotaxide</i> )
Pedicelli haud ita longi	...	29
29. Siliquæ longe rostratæ; rostrum valvarum trientem ad minimum æquans	...	30
Siliquæ multo brevius rostratæ vel subastylæ	...	33
30. Siliquæ pendulæ rostrum longissimum valvis plus duplo longius 5—6-spermum torulosum apice appланatum; caulis ramosissimus; folia parca, omnia petiolata; pedicelli flore breviores:		
		<i>B. longirostris Boiss.</i>
Siliquæ erectæ rostrum brevius, plerumque aspermum vel 1-spermum; pedicelli longiores	...	31
31. Rostrum valvis æquilongum subulatum; siliquæ 2—3-pollicares; folia caulina perpaucæ, summa squamæformia; pedicelli fructiferi longitudine valvarum	...	<i>B. oxyrrhina Coss.</i>
Rostrum crassius, obtusatum valvis triente vel dimidio brevius; pedicelli valvis breviores;	...	32
32. Folia caulina superiora exigua 3-angulari-squamæformia vel folia caulina fere omnia ad squamas reducta	...	<i>B. Tournefortii Gou.</i>
Caulis decrescenti-foliatus	...	<i>B. sabularia Brot.</i>
33. <i>Siliquæ tenuissime (capillari-) stipitatae</i> apice subtruncatæ, haud distincte stylatae, plano-compressæ, pendulæ	...	34
Siliquæ haud, vel crasse stipitatæ, distincte stylatae vel rostratæ, numquam pendulæ, teretusculæ vel subtetragonæ, rarius + compressæ	...	35
34. Petala calyce 2-plo longiora; folia latiuscula inciso-dentata vel pinnatifida	...	<i>B. Harra Forsk.</i>
(Diplotaxis pendula, D. crassifolia et D. Lagascana DC. & Pendulina Webbiana Willk.)		
Petala calyce paulo longiora; folia angusta spathulato-linearia indivisa vel lobulo lineari utrinque aucta	...	<i>B. intricata Willk.</i>
		(sub <i>Pendulina</i> .)
35. Stylus tenuis linearis vel basi constrictus obconicus a valvis manifeste separatus	...	36
Stylus robustior conicus, subulato-conicus vel subancipi compressus breviterque 3-angularis	...	43
36. Caulis brevissimus (2—3-pollicaris) aphyllus: petala calyce parum longiora; herba glabra annua	...	<i>B. scaposa DC.</i>
		(sub <i>Diplotaxide</i> )
Caulis + foliatus (rarissime aphyllus: in <i>B. murali</i> ); petala calyce 2-plo vel ultra longiora	...	37

37. Stylus basi constrictus obconicus (pedicelli sepalis æquilongi)  
*B. viminea Ardoino.*  
(Diplotaxis viminea DC.)
- Stylus linearis --- --- --- --- --- --- --- 38
38. Sepala patentia (caulis alte foliatus) --- --- --- --- --- --- 39  
Sepala erecta (caulis parte inferiore solum foliatus sub-  
scaposus) --- --- --- --- --- --- 41
39. Siliquæ pedicellis subæquilongæ (juniores alabastra superantes);  
flores lutei --- --- --- --- --- --- *B. tenuifolia Boiss.*  
(Diplotaxis tenuifolia DC.)
- Siliquæ pedicellis 2—3-plo longiores; flores albi vel carnei 40
40. Petala alba fusco venosa; caulis pilis rigidis patentibus usque  
ad inflorescentiam hispidus; folia lyrato-pinnati partita (stigma  
insigniter 2-lobum) --- --- --- --- --- *B. erucoides Ardoino.*  
(Diplotaxis erucoides DC. — D. hispidula Ten.)
- Petala pure carnea; folia sinuato-dentata; herba glabrescens  
*B. apula Ten.*  
(sub Diplotaxide)
41. Pedicelli calyce paulo breviores --- --- --- --- *B. monensis Huds.*  
Pedicelli calyces superantes --- --- --- --- 42
42. Sepala sparse hispido-pilosa; siliquæ juniores gemmas supe-  
rantes; annua --- --- --- --- --- *B. muralis Boiss.*  
(Diplotaxis muralis DC.)
- Sepala densius hispida; siliquæ juniores florum gemmis  
breviores; perennis --- --- --- --- --- *B. Gravinae Ten.*  
(B. nivalis Boiss. et Heldr. ?)
43. Folia caulina plane nulla vel subnulla, saltem valde diminuta  
basilaris difformia solumque ramos stipantia parva linearia --- 44  
Caulis continue decrescenti-foliatus --- --- --- --- 49
44. Pedicelli fere siliquæ dimidiæ æquantes --- --- --- --- 45  
Pedicelli siliqua multo breviores --- --- --- --- 46
45. Folia basilaria runcinato- et pectinato-pinnatisecta, lobis acu-  
minatis --- --- --- --- --- *B. lœvigata Lag.*  
Folia basilaria haud runcinata nec acuminato-lobata, sed  
lyrato-pinnatipartita --- --- --- --- --- *B. Cossoniana B. et R.*
46. Flores albi purpureo-venosi (siliquæ crassæ stylus longiusculus,  
calyx erectus) --- --- --- --- --- *B. insularis Mor.*  
Flores lutei concolores --- --- --- --- --- 47
47. Stylus subnnullus (stigma emarginatum); calyx demum de-  
flexus; siliquæ angustæ graciles --- --- --- --- *B. balearica Camb.*  
Stylus distinctus --- --- --- --- --- 48
48. Corymbus pauci-(1—6-)florus compactus; racemus fructifer

abbreviatus latus ambitu orbicularis vel ovalis; siliquæ latæ plerumque patentes	—	<i>B. humilis DC.</i> ( <i>B. saxatilis</i> et <i>B. repanda</i> DC.)
Corymbus 6—20-florus laxior; racemus fruetifer magis elongatus oblongus; siliquæ angustæ erectæ:	—	<i>B. Rouyanæ Janka</i>
( <i>Diplotaxis brassicoides</i> Rouy *) [ <i>Sinapis nūticaulis</i> Lag. — <i>B. Blancoana</i> <i>B. et R.</i> ]		
49. Pedicelli calyces manifeste superantes	—	50
Pedicelli calyces æquantes, breviores vel paulo tantum lon- giores	—	60
50. Folia omnia vel caulina inferiora lyrato-pinnatifida vel pinna- tipartita	—	51
Folia indivisa vel æqualiter sinuato-lobata	—	56
51. Stylus <i>tenuiter conicus</i> 2—3" longus, siliqua manifeste stipitata	—	52
Stylus <i>crasse conicus</i> 1—2" longus vel <i>applanato-triangu-</i> <i>laris</i> , siliqua haud vel substipitata	—	53
52. Flores magni racemosi; calyx cylindraceus accumbenti-erectus; siliquæ ultrapollicares	—	<i>B. Robertiana Gay</i>
Flores parvi corymbosi; calyx breviter aperte campanula- tus; siliquæ vix pollicares	—	<i>B. fruticulosa Cyr.</i>
53. Stylus <i>crasse conicus</i> 1—2" longus, siliqua substipitata:		
<i>B. rupestris Raf.</i>		
Stylus <i>applanato-3-angularis</i> , siliquæ haud stipitatae	—	54
54. Pedicelli infimi siliquis parum breviores; folia superiora sub- sessilia; herba parce retrorsum hirtula vel glabra (folia su- periora plerumque pinnatipartita, laciniis anguste linearibus):		
<i>B. catholica DC.</i>		
Pedicelli fructus $\frac{1}{2}$ v. $\frac{1}{3}$ partem æquantes; folia omnia petiolata; herba densius vestita	—	55
55. Inferne setoso-hispidissima; folia superiora sublinearia integra vel dentata	—	<i>B. virgata Cav.</i>
Retrorsum scabrido-pubescent; folia omnia pinnatipartita :		
<i>B. siifolia Kze.</i>		
56. Herbæ omni statu setuloso-scabrae	—	57
Herbæ glaberrimæ vel saltem statu evoluto glabrae læves	—	58
57. Siliquæ haud stipitatae; semina ovalia, compressa	—	<i>B. Erucastrum L.</i>
( <i>Erucastrum obtusangulum</i> Rehb., <i>Brassica obtusangula</i> Jess.)		
Siliquæ stipitatæ; semina globosa	—	<i>B. elongata Ehrh.</i>

\* Cfr. cl<sup>i</sup>. M. G. Rouy opusculem nuperrime emissum: «Étude des Diplotaxis européens de la section Brassicaria» in «Revue des Sciences Naturelles 1882»

58. Flores albi; pedicelli cum siliquis longiuscule stipitatis rectilinei (folia indivisa remote leviter crenulata vel denticulata):  
*B. armoracioides* Czern.  
 Flores flavescentes; siliquæ haud stipitatæ in pedicellis partulis erectæ... 59  
 59. Herba perennis, foliis sinuato-pectinato-lobatis; siliquæ valvæ 1-nerves... *B. palustris* Pirona.  
 Annua, folia integerrima vel dentata; siliquæ valvæ 3-nerves... *B. juncea* Czerniaëw.  
 (Conspicetus plantar. Charkov. et Ucran. pag. 7. n. 161; — Sinapis juncea L.— S. campestris Bess. & Brassica Besseriana Andrz. ex Trautvetter Enum. pl. songoric. a Dr. Schrenk collect. in «Bullet. soc. Mose.» a. 1869 I p. 134—135.)
60. Sepala erecta; folia omnia petiolata... 61  
 Sepala patentia; folia superiora sessilia... 64  
 61. Folia pleraque ± divisa... 62  
 Folia omnia indivisa, ad summum sinuata vel undulata (stigma emarginatum)... *B. Richeri* Vill.  
 62. Petala flava... 63  
 Petala alba, violaceo-venosa (folia pinnatisecta, siliquæ haud torulosæ)... *B. valentina* L.  
 63. Folia regulariter pinnatipartita, pedicelli breves crassi:  
*B. Cheiranthus* Vill.  
 Folia lyrata vel summa indivisa, pedicelli elongati tenues:  
*B. setigera* Gay.  
 64. Folia subæqualiter pinnatisecta (segmenta sinuato-dentata), superiora-linearia integerrima; pedicelli graciles siliquis sub-2-plo breviores (flores magni, pedicelli calyce haud breviores):  
*B. Pseudo-Sinapis* Lge.  
 Folia lyrata vel indivisa, superiora lanceolata vel oblonga dentata (rarius subintegra); pedicelli siliquis multo vel pluries breviores; flores medioeri vel parvi... 65  
 65. Folia pleraque indivisa; pedicelli breves *crassi* (calyce breviores, flores parvi); rostrum tetragono-compressum crassum pyramidale siliqua continuum... *B. Sinapis Vis.*  
 (B. Sinapistrum Boiss., Sinapis arvensis L., S. Schkuhriana Rehb.)  
 Folia lyrata; pedicelli graciles... 66  
 66. Folia superiora oblongo-linearia subintegra; flores parvi; rostrum *gracile* 2—5" longum aspermum; *foliorum caulinorum pinnae deflexae*... *B. Preslji* Janka.  
 (Sinapis virgata Pres.)

- Folia superiora lanceolata dentata; flores mediocri; rost-  
rum 2-spermum; foliorum pinnæ sursum patulæ ... *B. baetica* Boiss.  
(*Corynelobus baeticus* R. et Sch.)
67. Flores magni, petala violaceo-venosa; calyx elongatus cylin-  
draceus clausus basi subbisaccatus ... ... ... ... ... ... 68  
 Flores minores; petala avenia; calyx patulus (siliquæ bre-  
ves latæque oblongo-conicæ) ... ... ... ... ... ... 70
68. Pedicelli siliquarum dimidiatae aequantes; folia runcinata; calyx  
post anthesin auctus ... ... ... ... ... ... *B. vesicaria* L.  
 Pedicelli breviores; folia haud runcinata; calyx cito deci-  
duus immutatus ... ... ... ... ... ... 69
69. Rostrum valvas aequans vel superans; semina unilateraliter  
anguste diaphano-alata ... ... ... ... ... ... *B. Uechtritziana* Janka.  
(*Eruca longirostris* Uechtr.)
- Rostrum valvis brevius; semina exalata nec marginata: *B. Eruca* L.
70. Pedicelli (crassi) sepalis siliquisque breviores ... ... *B. hispida* Boiss.  
(*Sinapis hispida* Schousb.)
- Pedicelli sepalis siliquasque aequantes vel longiores ... ... 71
71. Sepala pedicellos aequantia; pedicelli rigidi siliquiscum sub-  
aequilongis patentissimi ... ... ... ... ... ... *B. alba* Boiss.  
(*Sinapis alba* L.)
- Sepala pedicellis breviores; siliquæ in pedicellis graciliori-  
bus arcuato-adscendentibus ... ... ... ... ... ... *B. dissecta* Boiss.  
(*Sinapis dissecta* Lag. — *S. ucranica* Czern.)
-