

Left-continuous t-norms in Fuzzy Logic: an Overview*

János Fodor

Dept. of Biomathematics and Informatics, Faculty of Veterinary Sci.
Szent István University, István u. 2, H-1078 Budapest, Hungary
E-mail: Fodor.Janos@aotk.szie.hu

Abstract: In this paper we summarize some fundamental results on left-continuous t-norms. First we study the nilpotent minimum and related operations in considerable details. This is the very first example of a left-continuous but not continuous t-norm in the literature. Then we recall some recent extensions and construction methods.

Keywords: Associative operations, Triangular norm, Residual implication, Left-continuous t-norm, Nilpotent minimum.

1 Introduction

The concept of Fuzzy Logic (FL) was invented by Lotfi Zadeh [20] and presented as a way of processing data by allowing partial set membership rather than only full or non-membership. This approach to set theory was not applied to engineering problems until the 70's due to insufficient small-computer capability prior to that time.

In the context of control problems (the most successful application area of FL), fuzzy logic is a problem-solving methodology that provides a simple way to arrive at a definite conclusion based upon vague, ambiguous, imprecise, noisy, or missing input information. FL incorporates a simple, rule-based "IF X AND Y THEN Z " approach to solving a control problem rather than attempting to model a system mathematically.

When one considers fuzzy subsets of a universe, in order to generalize the Boolean set-theoretical operations like intersection, union and complement, it is quite natural to use *interpretations* of logic connectives \wedge , \vee and \neg , respectively [12]. It is assumed that the conjunction \wedge is interpreted by a *triangular norm* (*t-norm* for short), the disjunction \vee is interpreted by a *triangular conorm* (shortly: *t-conorm*), and the negation \neg by a *strong negation*.

Although engineers have learned the basics of theoretical aspects of fuzzy sets and logic, from time to time it is necessary to summarize recent developments even in such a fundamental subject. This is the main aim of the present paper.

* This research has been supported in part by OTKA T046762.

Therefore, we focus on recent advances on an important and rather complex subclass of t-norms: on *left-continuous t-norms*. The standard example of a left-continuous t-norm is the *nilpotent minimum* [4,15]. Starting from our more than ten years old algebraic ideas, their elegant geometric interpretations make it possible to understand more on *left-continuous t-norms with strong induced negations*, and construct a wide family of them. Studies on properties of fuzzy logics based on left-continuous t-norms, and especially on the nilpotent minimum (NM) have started only recently; see [1,14,13,18,19] along this line.

2 Preliminaries

In this section we briefly recall some definitions and results will be used later. For more details see [5,12].

A bijection φ of the unit interval onto itself preserving natural ordering is called an *automorphism* of the unit interval. It is a continuous strictly increasing function satisfying boundary conditions $\varphi(0) = 0, \varphi(1) = 1$.

A *strong negation* N is defined as a strictly decreasing, continuous function $N: [0, 1] \rightarrow [0, 1]$ with boundary conditions $N(0) = 1, N(1) = 0$ such that N is involutive (i.e., $N(N(x)) = x$ holds for any $x \in [0, 1]$). A standard example of a strong negation is given by $N_{st}(x) = 1 - x$. Any strong negation N can be represented as a φ -*transform* of the standard negation (see [17])

$$N(x) = \varphi^{-1}(1 - \varphi(x))$$

for some automorphism φ of the unit interval. In this case the strong negation is denoted by N_φ .

A *t-norm* T is defined as a symmetric, associative and nondecreasing function $T: [0, 1]^2 \rightarrow [0, 1]$ satisfying boundary condition $T(1, x) = x$ for all $x \in [0, 1]$.

A *t-conorm* S is defined as a symmetric, associative and nondecreasing function $S: [0, 1]^2 \rightarrow [0, 1]$ satisfying boundary condition $S(0, x) = x$ for all $x \in [0, 1]$.

For any given t-norm T and strong negation N a function S defined by $S(x, y) = N(T(N(x), N(y)))$ is a t-conorm, called the *N -dual t-conorm of T* . In this case the triplet (T, S, N) is called a *De Morgan triplet*.

Well-accepted models for conjunction (AND), disjunction (OR), negation (NOT) are given by t-norms, t-conorms, strong negations, respectively. In this paper we will focus mainly on t-norms.

The definition of t-norms does not imply any kind of continuity. Nevertheless, such a property is desirable from theoretical as well as practical points of view.

A t-norm T is *continuous* if for all convergent sequences $\{x_n\}_{n \in \mathbb{N}}, \{y_n\}_{n \in \mathbb{N}}$ we have

$$T\left(\lim_{n \rightarrow \infty} x_n, \lim_{n \rightarrow \infty} y_n\right) = \lim_{n \rightarrow \infty} T(x_n, y_n).$$

The structure of continuous t-norms is well known, see [12] for more details, especially Section 3.3 on *ordinal sums*.

3 Left-continuous t-norms

In many cases, weaker forms of continuity are sufficient to consider. For t-norms, this property is *lower semicontinuity* [12, Section 1.3]. Since a t-norm T is non-decreasing and commutative, it is lower semicontinuous if and only if it is *left-continuous* in its first component. That is, if and only if for each $y \in [0, 1]$ and for all non-decreasing sequences $\{x_n\}_{n \in \mathbb{N}}$ we have

$$\lim_{n \rightarrow \infty} T(x_n, y) = T\left(\lim_{n \rightarrow \infty} x_n, y\right).$$

If T is a left-continuous t-norm, the operation $I_T : [0, 1]^2 \rightarrow [0, 1]$ defined by

$$I_T(x, y) = \sup\{t \in [0, 1] \mid T(x, t) \leq y\} \quad (1)$$

is called the *residual implication* (shortly: R-implication) generated by T . An equivalent formulation of left-continuity of T is given by the following property ($x, y, z \in [0, 1]$):

$$\mathbf{(R)} \quad T(x, y) \leq z \quad \text{if and only if} \quad I_T(x, z) \geq y.$$

We emphasize that the formula (1) can be computed for any t-norm T ; however, the resulting operation I_T satisfies condition (R) if and only if the t-norm T is left-continuous. An interesting underlying algebraic structure of left-continuous t-norms is a commutative, residuated integral l-monoid, see [6] for more details.

4 Nilpotent Minimum and Maximum

The first known example of a left-continuous but non-continuous t-norm is the so-called *nilpotent minimum* [4] denoted as $T^{\mathbf{nM}}$ and defined by

$$T^{\mathbf{nM}}(x, y) = \begin{cases} 0 & \text{if } x + y \leq 1, \\ \min(x, y) & \text{otherwise.} \end{cases} \quad (2)$$

It can be understood as follows. We start from a t-norm (the minimum), and re-define its value below and along the diagonal $\{(x, y) \in [0, 1] \mid x + y = 1\}$. So, the question is natural: if we consider any t-norm T and “annihilate” its original values below and along the mentioned diagonal, is the new operation always a t-norm? The general answer is “no” (although the contrary was “proved” in [15] where the same operation also appeared).

The definition (2) can be extended as follows. Suppose that φ is an automorphism of the unit interval. Define a binary operation on $[0, 1]$ by

$$T_{\varphi}^{\mathbf{nm}}(x, y) = \begin{cases} 0 & \text{if } \varphi(x) + \varphi(y) \leq 1 \\ \min(x, y) & \text{if } \varphi(x) + \varphi(y) > 1 \end{cases}. \quad (3)$$

Thus defined $T_{\varphi}^{\mathbf{nm}}$ is a t-norm and is called the φ -nilpotent minimum.

Clearly, the following equivalent form of $T_{\varphi}^{\mathbf{nm}}$ can be obtained by using the strong negation N_{φ} generated by φ :

$$T_{\varphi}^{\mathbf{nm}}(x, y) = \begin{cases} \min(x, y) & \text{if } y > N_{\varphi}(x) \\ 0 & \text{otherwise} \end{cases}.$$

Extension of $T_{\varphi}^{\mathbf{nm}}$ for more than two arguments is easily obtained and is given by $T_{\varphi}^{\mathbf{nm}}(x_1, \dots, x_n) = \min_{i=1, \dots, n} \{x_i\}$ if $\min_{i \neq j} \{\varphi(x_i) + \varphi(x_j)\} > 1$, and $T_{\varphi}^{\mathbf{nm}}(x_1, \dots, x_n) = 0$ otherwise.

The N_{φ} -dual t-conorm of $T_{\varphi}^{\mathbf{nm}}$, called the φ -nilpotent maximum, is defined by

$$S_{\varphi}^{\mathbf{nm}}(x, y) = \begin{cases} \max(x, y) & \text{if } \varphi(x) + \varphi(y) < 1 \\ 1 & \text{otherwise} \end{cases}.$$

Clearly, $(T_{\varphi}^{\mathbf{nm}}, S_{\varphi}^{\mathbf{nm}}, N_{\varphi})$ yields a De Morgan triple.

In the next theorem we list the most important properties of $T_{\varphi}^{\mathbf{nm}}$ and $S_{\varphi}^{\mathbf{nm}}$. These are easy to prove.

Theorem 1. *Suppose that φ is an automorphism of the unit interval. The t-norm $T_{\varphi}^{\mathbf{nm}}$ and the t-conorm $S_{\varphi}^{\mathbf{nm}}$ have the following properties:*

(a) *The law of contradiction holds for $T_{\varphi}^{\mathbf{nm}}$ as follows:*

$$T_{\varphi}^{\mathbf{nm}}(x, N_{\varphi}(x)) = 0 \quad \forall x \in [0, 1].$$

(b) *The law of excluded middle holds for $S_{\varphi}^{\mathbf{nm}}$:*

$$S_{\varphi}^{\mathbf{nm}}(x, N_{\varphi}(x)) = 1 \quad \forall x \in [0, 1].$$

(c) *There exists a number α_0 depending on φ such that $0 < \alpha_0 < 1$ and $T_{\varphi}^{\mathbf{nm}}$ is idempotent on the interval $]\alpha_0, 1]$:*

$$T_{\varphi}^{\mathbf{nm}}(x, x) = x \quad \forall x \in]\alpha_0, 1].$$

(d) *With the previously obtained α_0 , $S_{\varphi}^{\mathbf{nm}}$ is idempotent on the interval $[0, \alpha_0[$:*

$$S_{\varphi}^{\mathbf{nm}}(x, x) = x \quad \forall x \in [0, \alpha_0[.$$

(e) *There exists a subset X_{φ} of the unit square such that $(x, y) \in X_{\varphi}$ if and only if $(y, x) \in X_{\varphi}$ and the law of absorption holds on X_{φ} as follows:*

$$S_{\varphi}^{\mathbf{nm}}(x, T_{\varphi}^{\mathbf{nm}}(x, y)) = x \quad \forall (x, y) \in X_{\varphi}.$$

(f) There exists a subset Y_φ of the unit square such that $(x, y) \in Y_\varphi$ if and only if $(y, x) \in Y_\varphi$ and the law of absorption holds on Y_φ as follows:

$$T_\varphi^{\mathbf{nM}}(x, S_\varphi^{\mathbf{nM}}(x, y)) = x \quad \forall (x, y) \in Y_\varphi.$$

(g) If A, B are fuzzy subsets of the universe of discourse U and the α -cuts are denoted by A_α, B_α , respectively ($\alpha \in [0, 1]$), then we have

$$A_\alpha \cap B_\alpha = [T_\varphi^{\mathbf{nM}}(A, B)]_\alpha \quad \forall \alpha \in]\alpha_0, 1]$$

and

$$A_\alpha \cup B_\alpha = [S_\varphi^{\mathbf{nM}}(A, B)]_\alpha \quad \forall \alpha \in [0, \alpha_0[$$

where α_0 is given in (c).

(h) $T_\varphi^{\mathbf{nM}}$ is a left-continuous t-norm and $S_\varphi^{\mathbf{nM}}$ is a right-continuous t-conorm. \square

4.1 Where does Nilpotent Minimum Come from?

Nilpotent minimum has been discovered not by chance. There is a study on contrapositive symmetry of fuzzy implications [4]. A particular case of those investigations yielded nilpotent minimum. Some of the related results will be cited later in the present paper.

Let T be a left-continuous t-norm and N a strong negation. Consider the residual implication I_T generated by T , defined in (1).

Contrapositive symmetry of I_T with respect to N (CPS(N) for short) is a property that can be expressed by the following equality:

$$I_T(x, y) = I_T(N(y), N(x)) \quad \forall x, y \in [0, 1]. \quad (4)$$

Unfortunately, (4) is generally not satisfied for I_T generated by a left-continuous (even continuous) t-norm T . In [4] we proved the following result.

Theorem 2 ([4]). *Suppose that T is a t-norm such that condition **(R)** is satisfied, N is a strong negation. Then the following conditions are equivalent ($x, y, z \in [0, 1]$).*

- (a) I_T has contrapositive symmetry with respect to N ;
- (b) $I_T(x, y) = N(T(x, N(y)))$;
- (c) $T(x, y) \leq z$ if and only if $T(x, N(z)) \leq N(y)$.

In any of these cases we have

- (d) $N(x) = I_T(x, 0)$,
- (e) $T(x, y) = 0$ if and only if $x \leq N(y)$. \square

In the case of continuous t-norms we have the following unicity result (see also [11]).

Theorem 3. *Suppose that T is a continuous t-norm. Then I_T has contrapositive symmetry with respect to a strong negation N if and only if there exists an automorphism φ of the unit interval such that*

$$T(x, y) = \varphi^{-1}(\max\{\varphi(x) + \varphi(y) - 1, 0\}), \quad (5)$$

$$N(x) = \varphi^{-1}(1 - \varphi(x)). \quad (6)$$

In this case I_T is given by

$$I_T(x, y) = \varphi^{-1}(\min\{1 - \varphi(x) + \varphi(y), 1\}). \quad \square \quad (7)$$

When I_T is any R-implication and I_T does not have contrapositive symmetry then we can associate another implication with I_T . Suppose that T is a t-norm which satisfies condition **(R)**. Define a new implication associated with I_T as follows:

$$x \rightarrow_T y = \max\{I_T(x, y), I_T(N(y), N(x))\}. \quad (8)$$

If I_T has contrapositive symmetry then $x \rightarrow_T y = I_T(x, y) = I_T(N(y), N(x))$.

Define also a binary operation $*_T$ by

$$x *_T y = \min\{T(x, y), N[I_T(y, N(x))]\}. \quad (9)$$

Obviously, $*_T = T$ if (4) is satisfied by $I = I_T$. Even in the opposite case, this operation $*_T$ is a fuzzy conjunction in a broad sense and has several nice properties as we state in the next theorem.

Theorem 4. *Suppose that T is a t-norm such that **(R)** is true, N is a strong negation such that $N(x) \geq I_T(x, 0)$ for all $x \in [0, 1]$ and operations \rightarrow_T and $*_T$ are defined by (8) and (9), respectively. Then the following conditions are satisfied:*

- (a) $1 *_T y = y$;
- (b) $x *_T 1 = x$;
- (c) $*_T$ is nondecreasing in both arguments;
- (d) $x \rightarrow_T y \geq z$ if and only if $x *_T z \leq y$. \square

In Table 1 we list most common t-norms and corresponding operations I_T , $*_T$, \rightarrow_T , with $N(x) = 1 - x$.

Therefore, nilpotent minimum can be obtained as the conjunction $*_{\min}$. In general, $*_T$ is not a t-norm, not even commutative. Sufficient condition to assure that $*_T$ is a t-norm is given in the next theorem.

Theorem 5. *For a t-norm T and a strong negation N , if $y > N(x)$ implies $T(x, y) \leq N(I_T(y, N(x)))$ then $*_T$ is also a t-norm. \square*

T	$\min(x, y)$	$\max(x + y - 1, 0)$	xy
I_T	$1, x \leq y$ y otherwise	$\min(1 - x + y, 1)$	$\min 1, \frac{y}{x}$
$*_T$	$\min(x, y), x + y > 1$ $0, x + y \leq 1$	$\max(x + y - 1, 0)$	$\min xy, \frac{x + y - 1}{y}$
\rightarrow_T	$1, x \leq y$ $\max(1 - x, y), x > y$	$\min(1 - x + y, 1)$	$\max \frac{y}{x}, \frac{1 - x}{1 - y}$

Table 1. Some t-norms and associated connectives

4.2 Implications Defined by Nilpotent Minimum and Maximum

Consider the De Morgan triple $(T_\varphi^{\text{nm}}, S_\varphi^{\text{nm}}, N_\varphi)$ with an automorphism φ of the unit interval and define the corresponding S-implication:

$$I(x, y) = S_\varphi^{\text{nm}}(N_\varphi(x), y) \quad (10)$$

$$= \begin{cases} 1, & x \leq y \\ \max(N_\varphi(x), y), & x > y \end{cases} \quad (11)$$

One can easily prove that the R-implication defined by T_φ^{nm} coincides with the S-implication in (11).

Proposition 1. *Let φ be any automorphism of the unit interval. Then we have for all $x, y \in [0, 1]$ that*

$$I_{T_\varphi^{\text{nm}}}(x, y) = S_\varphi^{\text{nm}}(N_\varphi(x), y). \quad \square$$

As a trivial consequence, $I_{T_\varphi^{\text{nm}}}$ always has contrapositive symmetry with respect to N_φ .

Now we list the most important and attractive properties of $I_{T_\varphi^{\text{nm}}}$. Their richness is due to the fact that R- and S-implications coincide and thus advantageous features of both classes are combined.

1. $I_{T_\varphi^{\text{nm}}}(x, \cdot)$ is non-decreasing
2. $I_{T_\varphi^{\text{nm}}}(\cdot, y)$ is non-increasing
3. $I_{T_\varphi^{\text{nm}}}(1, y) = y$

4. $I_{T_\varphi^{\mathbf{nM}}}(0, y) = 1$
5. $I_{T_\varphi^{\mathbf{nM}}}(x, 1) = 1$
6. $I_{T_\varphi^{\mathbf{nM}}}(x, y) = 1$ if and only if $x \leq y$
7. $I_{T_\varphi^{\mathbf{nM}}}(x, y) = I_{T_\varphi^{\mathbf{nM}}}(N_\varphi(y), N_\varphi(x))$
8. $I_{T_\varphi^{\mathbf{nM}}}(x, 0) = N_\varphi(x)$
9. $I_{T_\varphi^{\mathbf{nM}}}(x, I_{T_\varphi^{\mathbf{nM}}}(y, x)) = 1$
10. $I_{T_\varphi^{\mathbf{nM}}}(x, \cdot)$ is right-continuous
11. $I_{T_\varphi^{\mathbf{nM}}}(x, x) = 1$
12. $I_{T_\varphi^{\mathbf{nM}}}(x, I_{T_\varphi^{\mathbf{nM}}}(y, z)) = I_{T_\varphi^{\mathbf{nM}}}(y, I_{T_\varphi^{\mathbf{nM}}}(x, z)) = I_{T_\varphi^{\mathbf{nM}}}(T_\varphi^{\mathbf{nM}}(x, y), z)$
13. $T_\varphi^{\mathbf{nM}}(x, I_{T_\varphi^{\mathbf{nM}}}(x, y)) \leq \min(x, y)$
14. $I_{T_\varphi^{\mathbf{nM}}}(x, y) \geq \min(x, y)$

Notice that $I_{T_\varphi^{\mathbf{nM}}}$ can also be viewed as a QL-implication defined by

$$\begin{aligned} S(x, y) &= S_\varphi^{\mathbf{nM}}(x, y), \\ N(x) &= N_\varphi(x) \\ T(x, y) &= \min(x, y) \end{aligned}$$

in (4), as one can check easily by simple calculus.

Therefore, this QL-implication (which is, in fact, an S-implication and an R-implication at the same time) also has contrapositive symmetry with respect to N_φ . Concerning this case, the following unicity result was proved in [4].

Theorem 6 ([4]). *Consider a QL-implication defined by $\max_\varphi(N_\varphi(x), T(x, y))$, where T is a t -norm. This implication has contrapositive symmetry with respect to N_φ if and only if $T = \min$. \square*

5 Extensions and Constructions

In this section we summarize some important results on left-continuous t -norms obtained by Jenei and other researchers.

5.1 Left-continuous t-norms with Strong Induced Negations

The notions and some of the results in the above Theorem 2 were formulated in a slightly more general framework in [7]. We restrict ourselves to the case of left-continuous t-norms with strong induced negations; i.e., T is a left-continuous t-norm and the function $N_T(x) = I_T(x, 0)$ (the negation induced by T) is a strong negation.

Moreover, in a sense, a converse statement of Theorem 2 was also established in [7]: If T is a left-continuous t-norm such that $N_T(x) = I_T(x, 0)$ is a strong negation, then (a), (b) and (c) necessarily hold with $N = N_T$.

Already in [3], we studied the above algebraic property (c). Geometric interpretations of properties (b) and (c) were given in [7] under the names of *rotation invariance* and *self-quasi inverse property*, respectively. More exactly, we have the following definition.

Definition 1. Let $T : [0, 1]^2 \rightarrow [0, 1]$ be a symmetric and non-decreasing function, and let N be a strong negation. We say that T admits the *rotation invariance* property with respect to N if for all $x, y, z \in [0, 1]$ we have

$$T(x, y) \leq z \quad \text{if and only if} \quad T(y, N(z)) \leq N(x).$$

In addition, suppose T is left-continuous. We say that T admits the *self quasi-inverse* property w.r.t. N if for all $x, y, z \in [0, 1]$ we have

$$I_T(x, y) = z \quad \text{if and only if} \quad T(x, N(y)) = N(z). \quad \square$$

For left-continuous t-norms, rotation invariance is exactly property (c) in Theorem 2, while self quasi-inverse property is just a slightly reformulated version of (b) there. Nevertheless, the following geometric interpretation was given in [7]. If N is the standard negation and we consider the transformation $\sigma : [0, 1]^3 \rightarrow [0, 1]^3$ defined by $\sigma(x, y, z) = (y, N(z), N(x))$, then it can be understood as a rotation of the unit cube with angle of $2\pi/3$ around the line connecting the points $(0, 0, 1)$ and $(1, 1, 0)$. Thus, the formula $T(x, y) \leq z \iff T(y, N(z)) \leq N(x)$ expresses that the part of the unit cube above the graph of T remains invariant under σ . This is illustrated in the first part of Figure 1.

The second part of Figure 1 is about the self quasi-inverse property which can be described as follows (for quasi-inverses of decreasing functions see [16]). For a left-continuous t-norm T , we define a function $f_x : [0, 1] \rightarrow [0, 1]$ as follows: $f_x(y) = N_T(T(x, y))$. It was proved in [7] that f_x is its own quasi-inverse if and only if T admits the self quasi-inverse property. Assume that N is the standard negation. Then the geometric interpretation of the negation is the reflection of the graph with respect to the line $y = 1/2$. Then, if it is applied to the partial mapping $T(x, \cdot)$, extend discontinuities of $T(x, \cdot)$ with vertical line segments. Then the obtained graph is invariant under the reflection with respect to the diagonal $\{(x, y) \in [0, 1] \mid x + y = 1\}$ of the unit square.




Fig. 1. Rotation invariance property (left). Self quasi-inverse property (right).

5.2 Rotation Construction

Theorem 7 ([9]). *Let N be a strong negation, t its unique fixed point and T be a left-continuous t-norm without zero divisors. Let T_1 be the linear transformation of T into $[t, 1]^2$. Let $I^+ =]t, 1]$, $I^- = [0, t]$, and define a function $T_{\text{rot}} : [0, 1]^2 \rightarrow [0, 1]$ by*

$$T_{\text{rot}}(x, y) = \begin{cases} T(x, y) & \text{if } x, y \in I^+, \\ N(I_{T_1}(x, N(y))) & \text{if } x \in I^+ \text{ and } y \in I^-, \\ N(I_{T_1}(y, N(x))) & \text{if } x \in I^- \text{ and } y \in I^+, \\ 0 & \text{if } x, y \in I^-. \end{cases}$$

Then T_{rot} is a left-continuous t-norm, and its induced negation is N .

When we start from the standard negation, the construction works as follows: take any left-continuous t-norm without zero divisors, scale it down to the square $[1/2, 1]^2$, and finally rotate it with angle of $2\pi/3$ in both directions around the line connecting the points $(0, 0, 1)$ and $(1, 1, 0)$. This is illustrated in Fig. 2.

Remark that there is another recent construction method of left-continuous t-norms (called rotation-annihilation) developed in [10].

5.3 Annihilation

Let N be a strong negation (i.e., an involutive order reversing bijection of the closed unit interval). Let T be a t-norm. Define a binary operation $T_{(N)} : [0, 1]^2 \rightarrow [0, 1]$ as follows:

$$T_{(N)}(x, y) = \begin{cases} T(x, y) & \text{if } x > N(y) \\ 0 & \text{otherwise.} \end{cases} \quad (12)$$




Fig. 2. T^{nM} as the rotation of the min, with the standard negation

We say that T can be N -annihilated when $T_{(N)}$ is also a t-norm. So, the question is: which t-norms can be N -annihilated? The above results show that $T = \min$ is a positive example.

A t-norm T is said to be a *trivial annihilation* (with respect to the strong negation N) if $N(x) = I_T(x, 0)$ holds for all $x \in [0, 1]$. It is easily seen that if a continuous t-norm T is a trivial annihilation then $T_{(N)} = T$.

Two t-norms T, T' are called N -similar if $T_{(N)} = T'_{(N)}$. Let T be a continuous non-Archimedean t-norm, and $\langle [a, b]; T_1 \rangle$ be a summand of T . We say that this summand is *in the center* (w.r.t. the strong negation N) if $a = N(b)$.

Theorem 8 ([8]). (a) Let T be a continuous Archimedean t-norm. Then $T_{(N)}$ is a t-norm if and only if $T(x, N(x)) = 0$ holds for all $x \in [0, 1]$.

(b) Let T be a continuous non-Archimedean t-norm. Then $T_{(N)}$ is a t-norm if and only if

- either T is N -similar to the minimum,
- or T is N -similar to a continuous t-norm which is defined by one trivial annihilation summand in the center. \square

Interestingly enough, the nilpotent minimum can be obtained as the limit of trivially annihilated continuous Archimedean t-norms, as the following result states.

Theorem 9 ([8]). There exists a sequence of continuous Archimedean t-norms T_k ($k = 1, 2, \dots$) such that

$$\lim_{k \rightarrow \infty} T_k(x, y) = T^{\text{nM}}(x, y) \quad (x, y \in [0, 1]).$$

Moreover, for all k , T_k is a trivial annihilation with respect to the standard negation.

The nilpotent minimum was slightly extended in [2] by allowing a weak negation instead of a strong one in the construction. Based on this extension, monoidal t-norm based logics (MTL) were studied also in [2], together with the involutive case (IMTL). Ordinal fuzzy logic, closely related to $T^{\mathbf{NM}}$, and its application to preference modelling was considered in [1]. Properties and applications of the $T^{\mathbf{NM}}$ -based implication (called R_0 implication there) were published in [14]. Linked to [2], the equivalence of IMTL logic and NM logic (i.e., nilpotent minimum based logic) was established in [13].

6 Conclusion

In this paper we have presented an overview of some fundamental results on left-continuous t-norms. The origin and basic properties of the very first left-continuous (and not continuous) t-norm called *nilpotent minimum* was recalled in some details. Extensions and general construction methods for left-continuous t-norms were also reviewed from the literature.

Acknowledgement

The author is grateful to S. Jenei for placing Figures 1–2 at his disposal.

References

1. B. De Baets, F. Esteva, J. Fodor and Ll. Godo, Systems of ordinal fuzzy logic with application to preference modelling, *Fuzzy Sets and Systems* **124** (2001) 353–359.
2. F. Esteva and L. Godo, Monoidal t-norm based logic: towards a logic for left-continuous t-norms, *Fuzzy Sets and Systems* **124** (2001) 271–288.
3. J.C. Fodor, A new look at fuzzy connectives, *Fuzzy Sets and Systems* **57** (1993) 141–148.
4. J.C. Fodor, Contrapositive symmetry of fuzzy implications, *Fuzzy Sets and Systems* **69** (1995) 141–156.
5. J.C. Fodor and M. Roubens, *Fuzzy Preference Modelling and Multicriteria Decision Support*, (Kluwer Academic Publishers, Dordrecht, 1994).
6. U. Höhle, Commutative, residuated l-monoids, in: U. Höhle and E.P. Klement, Eds., *Non-Classical Logics and their Applications to Fuzzy Subsets. A Handbook of the Mathematical Foundations of Fuzzy Set Theory*. (Kluwer Academic Publishers, Boston, 1995).
7. S. Jenei, Geometry of left-continuous t-norms with strong induced negations, *Belg. J. Oper. Res. Statist. Comput. Sci.* **38** (1998) 5–16.
8. S. Jenei, New family of triangular norms via contrapositive symmetrization of residuated implications, *Fuzzy Sets and Systems* **110** (2000) 157–174.
9. S. Jenei, Structure of left-continuous t-norms with strong induced negations. (I) Rotation construction, *J. Appl. Non-Classical Logics* **10** (2000) 83–92.

10. S. Jenei, Structure of left-continuous triangular norms with strong induced negations. (II) Rotation-annihilation construction, *J. Appl. Non-Classical Logics*, **11** (2001) 351–366.
11. P. Smets and P. Magrez, Implications in fuzzy logic, *Int. J. Approximate Reasoning* **1** (1987) 327–347.
12. E.P. Klement, R. Mesiar, and E. Pap, *Triangular Norms*, (Kluwer Academic Publishers, Dordrecht, 2000).
13. D. Pei, On equivalent forms of fuzzy logic systems NM and IMTL, *Fuzzy Sets and Systems* **138** (2003) 187-195.
14. D. Pei. R_0 implication: characteristics and applications, *Fuzzy Sets and Systems* **131** (2002) 297 - 302.
15. P. Perny, *Modélisation, agrégation et exploitation des préférences floues dans une problématique de rangement*, (PhD thesis, Université Paris-Dauphine, Paris, 1992).
16. B. Schweizer and A. Sklar, *Probabilistic Metric Spaces*, (North-Holland, New York, 1983).
17. E. Trillas, Sobre funciones de negación en la teoría de conjuntos difusos, *Stochastica* **III** (1979) 47–60.
18. S.-M. Wang, B.-S. Wang and F. Ren, NML, a schematic extension of F.Esteva and L.Godos logic MTL, *Fuzzy Sets and Systems* (to appear)
19. S.-M. Wang, B.-S. Wang and X.-Y. Wang, A characterization of truth-functions in the nilpotent minimum logic, *Fuzzy Sets and Systems* **145** (2004) 253-266.
20. L.A. Zadeh, Fuzzy sets, *Information and Control* **8** (1965) 338–353.